



Chloroplast Transition Metal Regulation for Efficient Photosynthesis

Schmidt, Sidsel Birkelund; Eisenhut, Marion; Schneider, Anja

Published in:
Trends in Plant Science

DOI:
[10.1016/j.tplants.2020.03.003](https://doi.org/10.1016/j.tplants.2020.03.003)

Publication date:
2020

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC-ND](#)

Citation for published version (APA):
Schmidt, S. B., Eisenhut, M., & Schneider, A. (2020). Chloroplast Transition Metal Regulation for Efficient Photosynthesis. *Trends in Plant Science*, 25(8), 817-828. <https://doi.org/10.1016/j.tplants.2020.03.003>

Review

Chloroplast Transition Metal Regulation for Efficient Photosynthesis

Sidsel Birkelund Schmidt,¹ Marion Eisenhut,^{2,*} and Anja Schneider^{3,*}

Plants require sunlight, water, CO₂, and essential nutrients to drive photosynthesis and fulfill their life cycle. The photosynthetic apparatus resides in chloroplasts and fundamentally relies on transition metals as catalysts and cofactors. Accordingly, chloroplasts are particularly rich in iron (Fe), manganese (Mn), and copper (Cu). Owing to their redox properties, those metals need to be carefully balanced within the cell. However, the regulation of transition metal homeostasis in chloroplasts is poorly understood. With the availability of the arabidopsis genome information and membrane protein databases, a wider catalogue for searching chloroplast metal transporters has considerably advanced the study of transition metal regulation. This review provides an updated overview of the chloroplast transition metal requirements and the transporters involved for efficient photosynthesis in higher plants.

Transition Metal Transporters Are Central to Chloroplast Metal Regulation and Photosynthesis

Chloroplasts harbor three types of membranes: a double (inner and outer) envelope and a thylakoid membrane. The envelope and thylakoid are separated by the stroma compartment. The membranes, and the transporters embedded therein, are selective barriers enabling exchange of ions and metabolites between the cytosol, the stroma, and the thylakoid lumen compartment. Chloroplast functioning and efficient photosynthesis for plant growth rely on the metal cofactor-mediated electron transport chain, in particular the transition metals Fe, Mn, and Cu [1]. However, as ions, their chemical properties can lead to generation of undesired reactive oxygen species (ROS) [2], which, together with their different binding affinities to proteins [3], greatly challenge the use of metals in oxygenic photosynthesis. To avoid harmful generation of ROS, plants chelate metal ions by organic ligands, such as nicotianamine, phenolic compounds, or organic acids (e.g., citrate, malate, ascorbate) [4]. Thus, the transport, homeostasis, and regulation of the individual transition metals are fundamental to optimizing chloroplast functioning.

The chloroplast contains more than 3000 proteins of which about 90 transporters are associated with the chloroplast envelope membranes, controlling the exchange of ions and metabolites [5,6]. However, most of the proteins involved in metal transport activities across the chloroplast membranes remain largely unknown. With the availability of arabidopsis (*Arabidopsis thaliana*) genome information [7] and establishment of various total cellular and more specific chloroplast protein membrane databases [8,9], a wider catalogue for searching new chloroplast metal transporters is now available. Moving forward, reverse genetic studies combined with targeted proteomics and bioinformatics approaches are likely to identify novel transporters [10,11]. This review provides a comprehensive description of chloroplast transition metal requirements, including an update on identified metal transport proteins for the distribution of Fe, Mn, and Cu within the chloroplast. The dynamics and mechanisms allowing the adaptability of the photosynthetic machinery to constantly changing metal concentrations are discussed.

Highlights

The transition metals Fe, Mn, and Cu have fundamental functions in the photosynthetic machinery but are also involved in undesired oxidative reactions when in excess. Thus, chloroplast transition metal concentrations need tight regulation to maintain efficient photosynthesis.

Chloroplast transition metal homeostasis is complex. It involves a sophisticated interplay of specific transport proteins, chaperones and carriers, and protein translocation systems.

The availability of full genomic information and organelle-specific protein databases of model plants, such as *A. thaliana*, allows the identification of chloroplast metal transport candidates. Using this information, the two transport proteins CMT1 and PAM71 were recently discovered to function in tandem for delivery of Mn to the oxygen-evolving complex in PSII.

¹Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, 1871 Frederiksberg C, Denmark

²Biochemie der Pflanzen, Heinrich-Heine-Universität Düsseldorf, 40225 Düsseldorf, Germany

³Molekularbiologie der Pflanzen (Botanik), Department Biologie I, Ludwig-Maximilians-Universität München, 82152 Martinsried, Germany

*Correspondence: m.eisenhut@hhu.de (M. Eisenhut) and anja.schneider@lrz.uni-muenchen.de (A. Schneider).



Chloroplast Transition Metal Requirements

Chloroplasts are rich in transition metals, with up to 80% of total leaf Fe and about 30% of leaf Cu allocated to the chloroplasts [12,13]. In contrast, intracellular Mn is mostly stored in vacuoles and, to a lesser extent, in chloroplasts [14]. The chloroplast metal concentration for Fe, Mn, and Cu is roughly 7 to 10 μg per 10^9 chloroplasts isolated from arabidopsis mesophyll cells, in a ratio of 5.8:1.0:0.5 (Fe:Mn:Cu) [15,16]. Within the chloroplasts, 60 to 80% of Fe, Mn, and Cu is found in thylakoids, reflecting the essential use of these metals in photosynthetic proteins and complexes [12,17], with a total number of 22 Fe atoms, four Mn atoms, and one Cu atom required per photosynthetic electron transport chain (Table 1).

Fe

Fe is the prominent transition metal for protein complexes in the photosynthetic electron chain (Table 1). It is present as Fe^{2+} ligated to PsbA and PsbD in photosystem II (PSII), in heme proteins of PSII and cytochrome b_6/f (Cyt b_6/f), and in Fe–Sulfur [Fe–S] clusters of Cyt b_6/f , photosystem I (PSI), and ferredoxin (Fd) (Figure 1, Key Figure, and Table 1). Biomass and seed yields in arabidopsis can be significantly increased through improved photosynthesis by overexpression

Table 1. Transition Metal Quota in Proteins of the Photosynthetic Electron Transport Chain

Metal	Chloroplast protein (Gene ID)	Ligand/cofactor	Function	Protein Data Bank in Europe	Refs
Mn	PSII PsbA and PsbC (AtCg00020 and AtCg00280)	Mn_4CaO_5	Water oxidation Electron transport	3jcu (spinach), 5xnm, 5xnl (pea)	[103] [104]
Fe	PSII PsbA and PsbD (AtCg00020 and AtCg00270) PsbE and PsbF ^a (AtCg00580 and AtCg00570)	Fe^{2+} heme	Electron transport ^e Photoprotection	5mdx (arabidopsis), 3jcu (spinach), 5xnm, 5xnl (pea)	[105] [103] [104]
Fe	Cyt b_6/f PetB (AtCg00720) ^b PetA (AtCg00540) ^c PetC (At4g03280) ^d	3 × heme heme [2Fe–2S]	Proton translocation Electron transport Electron transport	1q90 (<i>Chlamydomonas</i>), 4h44 (<i>Nostoc</i>)	[106] [107]
Fe	PSI PsaA and PsaB (AtCg00350 and AtCg00340) PsaC (AtCg01060)	1 × [4Fe–4S] 2 × [4Fe–4S]	Electron transport Electron transport	5zji (maize), 2wse (various plants), 5l8r (pea)	[108] [109] [110]
Fe	Fd Fd1 (At1g10960) Fd2 (At1g60950) Fd3 (At2g27510) Fd4 (At5g10000)	[2Fe–2S]	Electron transport	1pfd (parsley), 1gaq (maize)	[111] [112]
Cu	PC Pete1 (At1g76100) Pete2 (At1g20340)	$\text{Cu}^+/\text{Cu}^{2+}$	Electron transport	9pcy (bean)	[113]

^aHeme bound to PsbE and PsbF constitutes Cyt b_{556} a and b.

^bHeme bound to PetB constitutes Cyt b_6 .




^cHeme bound to PetA constitutes Cyt f .

^d[2Fe–2S] bound to PetC constitutes the Rieske FeS protein.

^e Fe^{2+} is involved in electron transport, despite no changes in its oxidation state.

Key Figure

Chloroplast Transition Metal Transport

(A) Iron (Fe)**(B) Manganese (Mn)****(C) Copper (Cu)**

Trends in Plant Science

(See figure legend at the bottom of the next page.)

of the Rieske [2Fe–2S] protein (PetC) [18], implying that Fe availability is not a bottleneck [18]. [Fe–S] clusters have an essential role in linear electron transport but also serve as redox cofactor of the NADH dehydrogenase-like complex (NDH) in cyclic electron flow around PSI [19], a mechanism to balance the levels of ATP and NADPH necessary for efficient photosynthesis. Fe also plays an essential auxiliary function as di-Fe cofactor in the enzyme plastid terminal oxidase (PTOX) [20,21], which is localized to the nonappressed regions of the thylakoids. PTOX protects the plastoquinone pool from overreduction by reducing the number of electrons available for photosynthetic reactions, specifically under abiotic stress [20,21].

In addition to the crucial role in photosynthesis, Fe is also involved as cofactor in many stroma and envelope localized processes. In these compartments, Fe-containing superoxide dismutase (FeSOD) [22] catalyzes the dismutation of harmful superoxide radicals (O_2^-) into hydrogen peroxide (H_2O_2), which is further decomposed by heme-containing ascorbate peroxidases (APX) to produce H_2O and O_2 [23]. Moreover, [Fe–S] clusters occur as cofactors of enzymes involved in chlorophyll metabolism, including chlorophyll *a* oxygenase (CAO), pheophorbide *a* oxygenase (PAO), 7-hydroxymethyl chlorophyll *a* reductase (HCAR) (for review, see [24]), and translocon at the inner chloroplast envelope 55 (TIC55) [25]. [Fe–S] clusters, together with siroheme (i.e., a specialized heme-like prosthetic group), are cofactors of sulfite and nitrite reductases (SiR and NiR) [26,27], both of which require reduced Fd as the physiological electron donor.

Mn

Mn is essential for light-driven oxidation of H_2O to extract the electrons needed in the photosynthetic electron chain and to release protons to generate the universal energy unit adenosine triphosphate (ATP). As a byproduct, O_2 is produced. For this purpose, four Mn ions constitute the catalytic center in the Mn_4CaO_5 cluster of the oxygen-evolving complex in PSII (Table 1 and Figure 1). In accordance with the Kok cycle [28], the Mn ions cycle through different oxidation states (Mn^{3+} , Mn^{4+}), the so-called S states, driven by the successive absorption of photons to extract electrons from H_2O [29]. PSII activity is therefore highly sensitive to Mn deficiency [30]. Moreover, a binuclear Mn^{2+} center is involved in recognition of the thylakoid-associated phosphatase 38 (TAP38) and phosphorylated light-harvesting chlorophyll *a/b* binding protein 1 (Lhcb1) during state transition; a mechanism allowing plants to regulate energy distribution between PSI and PSII [31,32].

Figure 1. (A) Iron (Fe) transport: At the outer envelope (OE) membrane, Fe uptake predominantly occurs as Fe-citrate. In the intermembrane space, ferric reductase oxidase 7 (FRO7) facilitates Fe reduction. Subsequently, Fe^{2+} is transported across the inner envelope membrane (IE) by the NiCo–PIC1 (nickel/cobalt transporter–permease in chloroplasts 1) transport system into the stroma, where it is provided for heme biogenesis and Fe–Sulfur [Fe–S] cluster assembly together with other Fe requiring enzymes (FeRE) or stored in ferritins when in excess. Frataxin (FH) delivers Fe to the [Fe–S] cluster assembly machinery, and [Fe–S] clusters are transferred to their target photosynthetic apoproteins via specific transfer proteins (TP), whereas the transport mechanisms for heme incorporation into photosystem II (PSII) and cytochrome (Cyt) *b₆/f* are unknown (grey arrows). Stromal Fe export is facilitated by the yellow stripe1-like (YSL)4/6 proteins. (B) Manganese (Mn) transport: The chloroplast Mn transporter 1 (CMT1) at the IE allows Mn uptake into the chloroplast stroma, where Mn^{2+} is required for activation of Mn responsive enzymes (MnRE). Stromal Mn^{2+} is imported into the thylakoid lumen via photosynthesis affected mutant 71 (PAM71), where it is assembled into the Mn_4CaO_5 cluster constituting the oxygen-evolving complex of PSII. The luminal protein PsbP possibly acts as a chaperone for Mn. (C) Copper (Cu) transport: The P-type ATPases, heavy metal ATPase (HMA)6 and HMA8, work in tandem for Cu^+ uptake at the IE and thylakoid membrane, respectively [40]. Furthermore, HMA1 functions in Cu uptake [and possibly export, depending on the actual situation (grey arrow)] at the IE. The plant-specific Cu chaperone plastid chaperone 1 (PCH1) provides the guided Cu^+ delivery to HMA6. The Cu chaperone for Cu/Zinc superoxide dismutase (Cu/ZnSOD) – CCS – is responsible for Cu delivery to Cu/ZnSOD and may also facilitate Cu delivery to HMA8 (grey arrow). The plastocyanin (PC) apoprotein is imported via the secretion system (Sec) pathway into the thylakoid lumen, where the protein is matured upon Cu incorporation. Abbreviations: Fd, ferredoxin; PetC, Rieske [2Fe–2S] protein; PSI, photosystem I; Tat, twin arginine translocase.

In the chloroplast stroma, enzymes of amino acid biosynthetic pathways are activated by Mn^{2+} . The imidazoleglycerol-phosphate dehydratase (IGPD) is a key enzyme in histidine biosynthesis [33–35]. The shikimate pathway of aromatic amino acid biosynthesis involves the Mn^{2+} -dependent 3-deoxy-D-*arabino*-heptulosonate 7-phosphate synthase (DAHP) enzyme [36]. Recently, a disputable hypothesis has been put forward, that Mn-binding to Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) favors its oxygenation activity [37]; by doing so, the oxygenase reaction of Rubisco would generate additional reductants, which participate in the assimilation of nitrate into amino acids to mitigate carbon/nitrogen imbalances derived from atmospheric CO_2 fluctuations.

Cu

Plastocyanin (PC) is the small soluble Cu protein in the thylakoid lumen, mediating the electron transfer from Cyt *b₆/f* to PSI (Table 1 and Figure 1). In *Arabidopsis*, two PC isoforms are encoded that have an equivalent essential function in electron transport activity. However, the PC1 isoform accumulates at lower levels than PC2, which is more sensitive to Cu availability at the protein level [38,39]. Another abundant Cu enzyme, polyphenol oxidase (PPO), contains two Cu^{2+}/Cu^+ ions per monomer. In the thylakoid lumen of spinach (*Spinacia oleracea*), the enzyme is thought to function in defense against biotic attacks [40]. However, genes encoding PPO are absent in *Arabidopsis* and the enzyme is therefore likely dispensable for the primary metabolism [41,42].

In the stroma compartment, the major Cu enzyme is the ROS-detoxifying Cu/Zinc superoxide dismutase (Cu/ZnSOD), where Cu^{2+}/Cu^+ acts as the redox active cofactor. Despite their colocalization in chloroplasts, FeSOD and Cu/ZnSOD share minimal structural homology and have evolved independently of one another [22,43]. The expression profiles of *SOD* genes in a number of plants indicate that both Cu/ZnSOD and FeSOD are even redundant under various conditions, such as salt, drought, and heat stress [44–46].

Chloroplast Responses to Environmental Transition Metal Deficiency

Transition metal deficiency manifests itself in reduced biomass production and a general discoloration of leaves, referred to as chlorosis. Cu deficient plants develop chlorotic symptoms that appear at the tip of the youngest leaves [47]. Fe and Mn deficiency cause interveinal chlorosis, which appears as a sharp distinction between veins and chlorotic areas of the youngest leaves [48,49].

At the cellular level, transition metal deficiency conditions often trigger disorganization of the chloroplast and its thylakoid membrane system. For instance, *Arabidopsis* permease in chloroplasts 1 (*pic1*) mutants are unable to import Fe into chloroplasts, and their chloroplast number and size are reduced. Chloroplast development was found to be distorted with inchoate or even absent thylakoids [50]. Likewise, in Mn-deficient chloroplasts of *Arabidopsis* chloroplast Mn transporter 1 (*cmt1*) mutants, chloroplast development was abnormal and thylakoids appeared disorganized, with either hypo- or hyper-stacked grana lamellae [15,16]. Clearly, Fe and Mn deficiencies directly affect the biogenesis of photosynthetic membrane complexes.

At the molecular level, genes associated with photosynthesis and tetrapyrrole metabolism are extensively downregulated under Fe deficiency [51]. Accordingly, impairment of chlorophyll biosynthesis is the most prominent sign of Fe deficiency. Decreased mRNA levels are also reflected in reduced protein levels of abundant Fe proteins, such as Fd (especially Fd2), proteins of the Cyt *b₆/f* complex, and enzymes of the S assimilation pathway [52]. Reduced accumulation of proteins of the [Fe–S] cluster assembly system, S mobilization A and B (SufA and SufB), presumably avoids the risk of forming incomplete [Fe–S] clusters, which are likely to accelerate the formation of the highly reactive hydroxyl radical in the Fenton reaction [52]. For the apoprotein SufA, the lack

of Fe as cofactor may cause protein instability rather than decreased mRNA levels [52]. Disrupting the transfer of [Fe-S] clusters to their target proteins, such as Fd, triggers leaf Fe-deficiency responses and, interestingly, results in overaccumulation of Fe in the chloroplast [53].

In the absence of Mn, the PSII supercomplexes are less stable [54,55]. A conceptual model explains the main events of Mn deficiency in PSII with Mn deficiency leading to reduced Mn binding in PSII complexes, which causes disintegration of PSII supercomplexes [56]. Consequently, the PSII core is increasingly affected by oxidative damage, which lowers PSII yield and eventually CO₂ assimilation and biomass production [30,54]. To prevent those detrimental effects, plants are able to mobilize stored Mn from the vacuole and recruit it to the chloroplast [57]. Recently, a group of ancient barley landraces was demonstrated to have a superior Mn efficiency trait, which enables the landraces to maintain Mn binding in PSII under Mn-deficient conditions [49].

Cu deficiency does not alter the expression level of the *PC* genes but instead triggers a large molecular remodeling that allows Cu to be preferentially allocated to PC [58–60]. In order to achieve this, transcripts of *Cu/ZnSOD* undergo microRNA-mediated downregulation. As a result, Cu provision is prioritized toward PC in plant chloroplasts [2,40,58,60,61].

Chloroplasts Membranes – Barriers to Transition Metal Import and Mobilization

Due to their endosymbiotic origin, chloroplasts are surrounded by a double-membrane envelope, and thus Fe, Mn, and Cu have to pass the outer envelope (OE) and the inner envelope (IE) (Figure 1). In addition, the thylakoid membrane system comprises the chloroplast internal extensive membrane system, and at least Mn²⁺ and Cu²⁺ need to cross this membrane to reach their destinations in the lumen (Figure 1).

Chloroplast Fe Transport

The uptake of Fe into chloroplasts is relatively well studied (Figure 1A). Studies in sugar beet (*Beta vulgaris*) [62] and oilseed rape (*Brassica napus* L.) [63] have demonstrated that Fe³⁺-citrate is the preferred substrate for chloroplast Fe uptake across the OE. Once in the chloroplast intermembrane space, Fe transport across the IE occurs most likely in the form of Fe²⁺ [64,65]. In Arabidopsis, it involves the ferric reductase oxidase 7 (FRO7) [66] facilitating Fe³⁺ reduction for the subsequent transport of Fe²⁺ across the IE. In sugar beet, the activity of the Fe³⁺ reductase enzyme was clearly associated with the IE and required the operation of the photosynthetic electron transport chain for providing reducing power [67].

PIC1 provided the first evidence for Fe²⁺ transport across the chloroplast IE membrane [50,64,68]. PIC1 is proposed to be part of a larger Fe import complex also involving the putative metal transport protein NiCo, which belongs to the Ni²⁺-Co²⁺ transporter family (TC#2.A.52) according to the Transporter Classification Database [69]. The current model is that Fe is first bound by NiCo and subsequently transferred to PIC1. The mode of Fe²⁺ uptake employs the proton motif force across the IE [64,68]. The strong coexpression of *PIC1*, *NiCo*, and *FRO7* observed in oilseed rape [63] supports the reduction-based Fe transport system of chloroplasts. An alternative pathway to PIC1-dependent Fe uptake has been suggested to involve the multiple antibiotic resistance 1 (MAR1) and the mitoferrin-like 1 (MFL1) proteins, although their integration into the IE have not been unequivocally shown. MFL1 belongs to the mitochondrial carrier family proteins but is annotated as an IE chloroplast protein [9]. It has a predicted chloroplast targeting peptide and might be involved in Fe transport under Fe excess [70]. Similarly, the MAR1 protein has been speculated to act as transporter for Fe chelators, such as citrate or nicotianamine [71]. However, the expression of *MAR1* in oilseed rape was much lower than the *FRO7-PIC1-NiCo* system under Fe-replete conditions [63] and is downregulated by 60% under Fe deficiency

[71], suggesting a minor importance of this transporter in Fe delivery under these conditions. Recently, a prokaryotic-type ATP-binding cassette (ABC) transporter module has been implicated in metal uptake at the IE membrane [72]. This module consists of the membrane intrinsic ABC subfamily I 12 (ABC112) and the soluble ABC110 component, which is the nucleotide-binding domain. A third player, ABC111, also known as NAP14 (for nonintrinsic ABC protein 14), might be attached to the IE and/or to plastoglobuli [72,73]. Analysis of *abc110* and *abc111* (*=nap14*) showed severe growth defects and strong chlorosis accompanied by impaired transition metal homeostasis in both mutants [72,73]. Whereas ABC111 (NAP14) is involved in Fe homeostasis [73], the transport specificity of the ABC10/ABC12 module at the IE awaits the molecular identification of the substrate binding component(s).

To regulate chloroplast Fe concentration, the two yellow stripe1-like (YSL) proteins YSL4 and YSL6, are hypothesized to function in Fe efflux from the chloroplast to the cytoplasm. The envelope localization of the proteins is still under debate, but the expression of the genes is upregulated in response to Fe excess [74]. In *Zea mays*, the Fe deficiency-related 4 (ZmFDR4) protein, a probable thylakoid membrane-localized protein, is more strongly expressed under Fe deficient than under Fe replete conditions [75]. Thus, FDR4 could be involved in Fe uptake across the thylakoid membrane in monocots.

Chloroplast Mn Transport

Despite the fundamental need for Mn in oxygenic photosynthesis, chloroplast Mn transporters were only recently identified (Figure 1B). The IE CMT1 [15,16] and the thylakoid photosynthesis affected mutant 71 (PAM71) protein [54], synonymous to putative chloroplast calcium (Ca) proton antiporter (CCHA) [76], act in tandem to efficiently deliver Mn to the thylakoid lumen for its incorporation into the Mn_4CaO_5 cluster in PSII [15,77]. CMT1 and PAM71 belong to the unknown protein family UPF0016 [78]. This protein family was recently established as functioning in transport processes. Loss of function *cmt1* arabidopsis mutants show strongly reduced growth accompanied by chloroplasts having aberrant thylakoid membrane systems and diminished integral photosynthesis protein complexes due to strongly reduced chloroplast Mn content. Plant growth can hardly be complemented by supplementation of excess Mn concentrations in the growth medium, and CMT1 is therefore likely to be the dominant Mn uptake system at the IE of chloroplasts [15,16]. The *pam71* mutant grows comparably better than the *cmt1* mutant and is specifically impaired in PSII activity. This is caused by reduced Mn incorporation into PSII supercomplexes [54]. Notably, loss of PAM71 in both arabidopsis and the green alga *Chlamydomonas reinhardtii* can be compensated by enhanced Mn additions, indicating the presence of an uncharacterized low-affinity uptake system for Mn at the thylakoid membrane [54,78].

For the CMT1 and PAM71 proteins, an additional role in Ca transport has been suggested, and the proteins were named bivalent cation transporter (BICAT) 1 and 2, respectively [79]. BICAT1/PAM71 is proposed to transport Ca^{2+} into the thylakoid lumen and BICAT2/CMT1 to transport Ca^{2+} across the chloroplast IE [79]. However, Ca contents do not differ between *cmt1* mutant chloroplasts and wild-type (WT) chloroplasts [16], and *pam71* mutants even accumulate more Ca in the thylakoid lumen with respect to the WT [54]. The physiological effects in cyanobacteria mutants of the orthologous gene [80,81] underline the main function of the UPF0016 family is Mn transport in oxygenic photosynthetic organisms.

Chloroplast Cu Transport

The chloroplast Cu uptake and intraorganellar distribution is facilitated by two $\text{P}_{1\text{B}}$ -type ATPases, P-type ATPase of arabidopsis 1 and 2 (PAA1 and PAA2), also known as heavy

metal ATPase 6 and 8 (HMA6 and HMA8), which function in tandem in the transport of mono-valent Cu^+ ions across the chloroplast envelope and the thylakoid membrane, respectively [13,82,83] (Figure 1C). Biochemical analyses have revealed that both proteins are high-affinity Cu^+ transporters but differ in their biochemical properties as HMA8 has a higher affinity for Cu^+ than HMA6 [84]. Cu excess in the stroma triggers HMA8 degradation [59]. The phenotypes of the arabidopsis mutants *hma6* and *hma8* specifically relate to reduced Cu delivery to the major plastid Cu sinks. As expected, the *hma6* mutant shows impaired stromal Cu/ZnSOD activity, and both *hma6* and *hma8* mutants have a strong reduction in holo-PC protein content, resulting in impaired photosynthetic electron transport [82]. An additional member of the HMA family, the broad-specificity HMA1 protein, is suggested to facilitate ATP-dependent Cu import at the chloroplast IE. This pathway is distinct from HMA6 and becomes vital only under high-light conditions [85,86]. Besides Cu import into arabidopsis chloroplasts, Cu (and Zn) export activity has been reported for HvHMA1 in barley chloroplasts for remobilization purposes [87,88].

Transport Pathways and Mechanisms Ensuring Correct Protein Metalation in Chloroplasts

Chloroplast metalloproteins functioning in photosynthetic electron transport are assumed to obtain their metal from exchangeable stromal pools and are subject to the universal order of protein divalent metal-binding affinities described by Irving and Williams [3], with Cu forming tighter complexes than Fe and Mn, respectively. This challenges the use of transition metals in photosynthesis to ensure correct metal acquisition by apometalloproteins but can be overcome by either compartmentalized protein folding or by assistance of metal delivery proteins, known as metallochaperones. Metallochaperones pass metals between transporters and their target apoproteins via ligand-exchange reactions. Thus, they participate in the final transport step before crossing the thylakoid membrane, which is mediated by one of two distinct pathways; the chloroplast secretion system (Sec) or the chloroplast twin arginine translocase (Tat) pathway [89]. The translocon of the Sec-type pathway only accepts unfolded proteins, whereas the Tat-type pathway accommodates transport of mature and highly folded proteins [90].

Fe Metalation

Assembly of [Fe-S] clusters needed in stroma-localized proteins and proteins of the electron transport chain takes place in the stroma compartment by the SUF pathway. In the first stage, [Fe-S] clusters are built from Fe and S delivered by proteins onto so-called scaffold proteins, which are the primary sites of *de novo* cluster assembly [24]. However, the source of Fe and how Fe is delivered to those scaffold proteins are unknown. The Fe-storage protein ferritin (FER) was considered a candidate; however, the triple *fer1fer3fer4* arabidopsis mutant of three major FERs retained fully functional photosynthetic apparatus, suggesting that FERs are not the major Fe pool for the SUF pathway [91]. Instead, current models assume that a metallochaperone acquires Fe and directly donates Fe to the assembly pathway by interacting with one of the scaffold proteins. Frataxin (FH) is a metallochaperone candidate for [Fe-S] cluster assembly [92,93] (Figure 1). After assembly, [Fe-S] clusters are transferred to their respective target recipient apoprotein with the help of transfer proteins. A total of nine potential chloroplastic [Fe-S] transfer proteins have been identified so far [24]. The unidirectional and intact transfer of [Fe-S] clusters from one transfer protein to another indicates that some [Fe-S] transfer proteins work in sequence and most possess specificity to individual [Fe-S] cluster types. For example, [4Fe-4S] of PSI proteins are supplied by the chloroplast-localized nifU-like proteins NFU2 and NFU3, and high-chlorophyll fluorescence 101 (HCF101) [24,94,95]. The Asn-Glu-Glu-Thr motif protein NEET is proposed to transfer [2Fe-2S] clusters to Fd [53]. It remains elusive which transfer

protein delivers the [2Fe–2S] cluster to PetC (Table 1) prior to translocation across the thylakoid membrane via the Tat pathway to reach its destination on the luminal side of the Cyt *b₆/f* complex [96] (Figure 1).

Mn Metalation

The three luminal proteins PsbO, PsbP, and PsbQ of PSII are known to protect and stabilize the oxygen-evolving complex and maintain optimal Ca and chloride concentrations in plant chloroplasts [97]. The crystal structure of PsbP from spinach was shown to contain two specific Mn²⁺-binding sites, with one site having a relatively low affinity [98], making it a suitable candidate site for Mn donation to the Mn₄CaO₅ cluster assembly. Thus, a role for PsbP as a metallochaperone for Mn²⁺ in the thylakoid lumen has been suggested [56,98–100]. In line with this assumption, the ancestral form of PsbP in cyanobacteria, CyanoP, has been proposed as a promising Mn chaperone. Experimental support derives from the observation that the *Synechocystis* Δ*SynPAM71* mutant overaccumulates CyanoP, maybe to buffer excess amounts of Mn in this strain [81].

Cu Metalation

The plant-specific Cu chaperone plastid chaperone 1 (PCH1), generated by an alternative splicing event of *HMA6* pre-mRNA, resides in the intermembrane space. PCH1 contains an N-terminal metal-binding domain, which enables transient interaction of PCH1 to the intramembrane metal-binding site of HMA6 for Cu delivery across the IE [40,61,101]. In the stroma, another Cu chaperone, Cu chaperone for Cu/ZnSOD (CCS), is responsible for Cu delivery to Cu/ZnSOD [102]. CCS was further identified as a possible source for the targeted Cu provision to the thylakoid transporter HMA8. Thus, CCS might be involved in Cu delivery to PC [101] (Figure 1). Even so, the CCS knockout mutant did not show alterations in PSII activity, nor changed PSI and PSII protein amounts, even after high-light stress, when grown on high Cu media [102]. These findings imply the existence of further Cu metallochaperones in the chloroplast stroma. PC is a Sec substrate, and thus Cu insertion and PC maturation must be completed within the thylakoid lumen, suggesting that Cu chaperones are also required in this compartment.

Concluding Remarks and Future Perspectives

Chloroplasts respond instantaneously to changes in transition metal concentrations to regulate the mechanisms controlling transition metal transport and storage in the cell. Despite recent progress in identification and characterization of chloroplast transition metal transporters, major questions await answering, including the exact transport mode, the involvement of chaperones, and the nature of their substrate state. For the proteins involved, the unequivocal determination of their subcellular localization is essential. This is of special importance as accumulating evidence points towards dual organelle functions (in, e.g., chloroplast and mitochondrion/vacuole/endoplasmic reticulum) for a subset of transport proteins involved.

Although fundamental, little is known about chloroplast metal sensing (Box 1) and how metalloproteins can discriminate and acquire the correct metal cofactor in the complex matrix of the intermembrane space, stroma, and lumen fractions. This adds an additional layer of complexity to chloroplast transition metal regulation. Future work should integrate molecular metal sensors and omics approaches to gain a comprehensive knowledge of the sophisticated networks of response towards depletion and repletion of transition metal ions. This will provide new knowledge of chloroplast transition metal regulation in response to fluctuating transition metal availability and will contribute to our understanding and ultimate engineering of plant photosynthetic efficiency and productivity (see Outstanding Questions).

Outstanding Questions

How is the demand of the individual transition metals coordinated?

How is sequestration and remobilization of transition metals into/from storage pools (e.g., the vacuole) regulated?

How are heme cofactors assembled into photosynthetic proteins?

What is the transport mode of the UPF0016 family proteins CMT1 and PAM71 driving Mn transport?

Are Mn chaperones present in the chloroplast stroma and thylakoid lumen?

What is the molecular nature of the uncharacterized low-affinity Mn uptake systems at the thylakoid membrane and/or chloroplast envelope membrane? Do these low affinity systems also accept other transition metals, such as Cu and Fe?

Does a further stromal Cu chaperone exist for the thylakoid membrane-localized HMA8 transporter?

Are there specific ligands for Mn, Fe, or Cu in the chloroplast or are ligands similar to metal–ligand complexes in phloem and xylem sap?

Box 1. Chloroplast Metal Sensing and Regulation

When exposed to either metal deficiency or excess, the chloroplast needs to react instantaneously to avoid harmful imbalances affecting the photosynthetic electron transfer chain. Despite the central importance of maintaining photosynthetic efficiency, the knowledge on metal orchestration, metal interactions, and signaling within the chloroplast is surprisingly poor.

The main questions of chloroplast transition metal regulation are how metal concentrations and individual ratios are sensed within the chloroplast and how the specific metal concentration is eventually controlled. Impaired electron transfer, owing to imbalanced availability of metalloproteins of the photosynthesis apparatus, leads to generation of overreduction signaling molecules, such as ROS [15]. It is well known that information via ROS serves to tune processes inside and outside the chloroplast [114,115]. Metal homeostasis strategies, such as metal chelation and transport, are under nuclear transcriptional control and induced upon transition metal deficiency or excess [40,51,116]. Although speculative, the chloroplast may act as a cellular sensing and signaling site for metal deficiencies, and ROS production might induce a signaling cascade, regulating the expression of genes coding for metal importers (or exporters) at the IE. The analysis of the *cmt1* mutant supports the central role of metal transporters at the IE in chloroplast metal regulation; loss of the Mn transporter CMT1 results in strongly reduced Mn concentrations in the chloroplast leading to reduced PSII activity [15,16]. To accommodate deficiencies in PSII, adjustment of photosystem stoichiometry is achieved by decreasing subunits of PSI and Cyt *b₆/f*. Hence, surplus Fe has to be sequestered. Possibly, excess Fe is exported from the chloroplast and sequestered into the vacuole, the central cellular hub for transition metals. Thus, low Mn concentration implicates reduced levels of chloroplast Fe, yet Cu concentrations are unaffected [15].

Not only that, the chloroplast sends signals to the nucleus (retrograde signaling), cytosolic metal sensing triggers anterograde pathway (e.g., when plants experience Mn stress, expression of *CMT1* is downregulated) [15], and excess cytosolic Fe triggers *FER* expression [116]. Chloroplast transition metal requirement is dynamically changing and thus requires not only communication between this organelle and the nucleus but also long-distance signaling between the shoot and the root [117] as primary source for transition metal acquisition.

Acknowledgments

This research was financially supported by the Independent Research Fund Denmark – Technology and Production Sciences (grant DFF-5054-00042 to S.B.S) and the German Research Foundation (Deutsche Forschungsgemeinschaft) – EI945/3-1 to M.E. and SCHN560/4-1 to A.S. The authors wish to thank Debbie Maizels, Zoobotanica Scientific Illustration, for providing illustrations for Figure 1.

References

1. Merchant, S. (2006) Trace metal utilization in chloroplasts. In *The Structure and Function of Plastids* (Wise, R. and Hooper, J.K., eds), pp. 199–218, Springer
2. Ravet, K. and Pilon, M. (2013) Copper and iron homeostasis in plants: the challenges of oxidative stress. *Antioxid. Redox Signal.* 19, 919–932
3. Irving, H. and Williams, R.J.P. (1948) Order of stability of metal complexes. *Nature* 162, 746–747
4. Bashir, K. *et al.* (2016) Regulating subcellular metal homeostasis: the key to crop improvement. *Front. Plant Sci.* 7, 1192
5. Bouchnak, I. *et al.* (2019) Unravelling hidden components of the chloroplast envelope proteome: opportunities and limits of better MS sensitivity. *Mol. Cell. Proteomics* 18, 1285–1306
6. Finazzi, G. *et al.* (2015) Ion channels/transporters and chloroplast regulation. *Cell Calcium* 58, 86–97
7. The Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408, 796–815
8. Schwacke, R. *et al.* (2003) ARAMEMNON, a novel database for Arabidopsis integral membrane proteins. *Plant Physiol.* 131, 16–26
9. Ferro, M. *et al.* (2010) AT_CHLORO, a comprehensive chloroplast proteome database with subplastidial localization and curated information on envelope proteins. *Mol. Cell. Proteomics* 9, 1063–1084
10. Rolland, N. *et al.* (2012) The biosynthetic capacities of the plastids and integration between cytoplasmic and chloroplast processes. *Annu. Rev. Genet.* 46, 233–264
11. Ling, Q. *et al.* (2019) Ubiquitin-dependent chloroplast-associated protein degradation in plants. *Science* 363, eaav4467
12. Terry, N. and Low, G. (1982) Leaf chlorophyll content and its relation to the intracellular-localization of iron. *J. Plant Nutr.* 5, 301–310
13. Shikanai, T. *et al.* (2003) PAA1, a P-type ATPase of *Arabidopsis*, functions in copper transport in chloroplasts. *Plant Cell* 15, 1333–1346
14. Pittman, J.K. (2005) Managing the manganese: molecular mechanisms of manganese transport and homeostasis. *New Phytol.* 167, 733–742
15. Eisenhut, M. *et al.* (2018) The plastid envelope CHLOROPLAST MANGANESE TRANSPORTER1 is essential for manganese homeostasis in *Arabidopsis*. *Mol. Plant* 11, 955–969
16. Zhang, B. *et al.* (2018) Inner envelope CHLOROPLAST MANGANESE TRANSPORTER 1 supports manganese homeostasis and phototrophic growth in *Arabidopsis*. *Mol. Plant* 11, 943–954
17. Lysenko, E.A. *et al.* (2019) Distribution of Cd and other cations between the stroma and thylakoids: a quantitative approach to the search for Cd targets in chloroplasts. *Photosynth. Res.* 139, 337–358
18. Simkin, A.J. *et al.* (2017) Overexpression of the RieskeFeS protein increases electron transport rates and biomass yield. *Plant Physiol.* 175, 134–145
19. Laughlin, T.G. *et al.* (2019) Structure of the complex I-like molecule NDH of oxygenic photosynthesis. *Nature* 566, 411–414
20. Nawrocki, W.J. *et al.* (2015) The plastid terminal oxidase: its elusive function points to multiple contributions to plastid physiology. *Annu. Rev. Plant Biol.* 66, 49–74
21. Krieger-Liszka, A. and Feilke, K. (2015) The dual role of the plastid terminal oxidase PTOX: between a protective and a pro-oxidant function. *Front. Plant Sci.* 6, 1147

22. Pilon, M. *et al.* (2011) The biogenesis and physiological function of chloroplast superoxide dismutases. *Biochim. Biophys. Acta* 1807, 989–998
23. Wada, K. *et al.* (2003) Crystal structure of chloroplastic ascorbate peroxidase from tobacco plants and structural insights into its instability. *J. Biochem.* 134, 239–244
24. Lu, Y. (2018) Assembly and transfer of iron–sulfur clusters in the plastid. *Front. Plant Sci.* 9, 336–336
25. Hauenstein, M. *et al.* (2016) A role for TIC55 as a hydroxylase of phytylcholins, the products of chlorophyll breakdown during plant senescence. *Plant Cell* 28, 2510–2527
26. Nakano, S. *et al.* (2012) The reductive reaction mechanism of tobacco nitrite reductase derived from a combination of crystal structures and ultraviolet–visible microspectroscopy. *Proteins* 80, 2035–2045
27. Kim, J.Y. *et al.* (2016) Structural and mutational studies of an electron transfer complex of maize sulfite reductase and ferredoxin. *J. Biochem.* 160, 101–109
28. Kok, B. *et al.* (1970) Cooperation of charges in photosynthetic O₂ evolution-I. A linear four step mechanism. *Photochem. Photobiol.* 11, 457–475
29. Shen, J.-R. (2015) The structure of photosystem II and the mechanism of water oxidation in photosynthesis. *Annu. Rev. Plant Biol.* 66, 23–48
30. Schmidt, S.B. *et al.* (2016) Photosystem II functionality in barley responds dynamically to changes in leaf manganese status. *Front. Plant Sci.* 7, 1–12
31. Pribil, M. *et al.* (2010) Role of plastid protein phosphatase TAP38 in LHClI dephosphorylation and thylakoid electron flow. *PLoS Biol.* 8, e1000288
32. Wei, X. *et al.* (2015) Structural mechanism underlying the specific recognition between the *Arabidopsis* state-transition phosphatase TAP38/PPH1 and phosphorylated light-harvesting complex protein Lhcb1. *Plant Cell* 27, 1113–1127
33. Ingle, R.A. (2011) Histidine biosynthesis. *Arabidopsis Book* 9, e0141
34. Bisson, C. *et al.* (2015) Crystal structures reveal that the reaction mechanism of imidazoleglycerol-phosphate dehydratase is controlled by switching Mn(II) coordination. *Structure* 23, 1236–1245
35. Rawson, S. *et al.* (2018) Elucidating the structural basis for differing enzyme inhibitor potency by cryo-EM. *Proc. Natl. Acad. Sci. U. S. A.* 115, 1795–1800
36. Entus, R. *et al.* (2002) Redox regulation of *Arabidopsis* 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase. *Plant Physiol.* 129, 1866–1871
37. Bloom, A.J. and Lancaster, K.M. (2018) Manganese binding to Rubisco could drive a photorespiratory pathway that increases the energy efficiency of photosynthesis. *Nat. Plants* 4, 414–422
38. Abdel-Ghany, S.E. (2009) Contribution of plastocyanin isoforms to photosynthesis and copper homeostasis in *Arabidopsis thaliana* grown at different copper regimes. *Planta* 229, 767–779
39. Pesaresi, P. *et al.* (2009) Mutants, overexpressors, and interactors of *Arabidopsis* plastocyanin isoforms: revised roles of plastocyanin in photosynthetic electron flow and thylakoid redox state. *Mol. Plant* 2, 236–248
40. Aguirre, G. and Pilon, M. (2016) Copper delivery to chloroplast proteins and its regulation. *Front. Plant Sci.* 6, 1250
41. Schubert, M. *et al.* (2002) Proteome map of the chloroplast lumen of *Arabidopsis thaliana*. *J. Biol. Chem.* 277, 8354–8365
42. Tran, L.T. *et al.* (2012) The polyphenol oxidase gene family in land plants: lineage-specific duplication and expansion. *BMC Genomics* 13, 395
43. Case, A.J. (2017) On the origin of superoxide dismutase: an evolutionary perspective of superoxide-mediated redox signaling. *Antioxidants (Basel)* 6, E82
44. Feng, K. *et al.* (2016) The SOD gene family in tomato: identification, phylogenetic relationships, and expression patterns. *Front. Plant Sci.* 7, 1279
45. Feng, X. *et al.* (2015) Genome-wide identification and characterization of the superoxide dismutase gene family in *Musa acuminata* cv. Tianbaojiao (AAA group). *BMC Genomics* 16, 823
46. Verma, D. *et al.* (2019) Genome-wide identification and characterization of abiotic-stress responsive SOD (superoxide dismutase) gene family in *Brassica juncea* and *B. rapa*. *BMC Genomics* 20, 227
47. Printz, B. *et al.* (2016) Copper trafficking in plants and its implication on cell wall dynamics. *Front. Plant Sci.* 7, 601
48. Briat, J.-F. *et al.* (2015) Iron nutrition, biomass production, and plant product quality. *Trends Plant Sci.* 20, 33–40
49. Schmidt, S.B. *et al.* (2019) Ancient barley landraces adapted to marginal soils demonstrate exceptional tolerance to manganese limitation. *Ann. Bot.* 123, 831–843
50. Duy, D. *et al.* (2007) PIC1, an ancient permease in *Arabidopsis* chloroplasts, mediates iron transport. *Plant Cell* 19, 986–1006
51. Rodríguez-Celma, J. *et al.* (2013) The transcriptional response of *Arabidopsis* leaves to Fe deficiency. *Front. Plant Sci.* 4, 276
52. Hantzis, L.J. *et al.* (2018) A program for iron economy during deficiency targets specific Fe proteins. *Plant Physiol.* 176, 596–610
53. Zandalinas, S.I. *et al.* (2020) Expression of a dominant-negative AtNEET-H89C protein disrupts iron-sulfur metabolism and iron homeostasis in *Arabidopsis*. *Plant J.* 101, 1152–1169
54. Schneider, A. *et al.* (2016) The evolutionarily conserved protein PHOTOSYNTHESIS AFFECTED MUTANT71 is required for efficient manganese uptake at the thylakoid membrane in *Arabidopsis*. *Plant Cell* 28, 892–910
55. Schmidt, S.B. *et al.* (2015) Metal binding in photosystem II super- and subcomplexes from barley thylakoids. *Plant Physiol.* 168, 1490–1502
56. Schmidt, S.B. *et al.* (2016) Manganese deficiency in plants: the impact on photosystem II. *Trends Plant Sci.* 21, 622–632
57. Lanquar, V. *et al.* (2010) Export of vacuolar manganese by AtNRAMP3 and AtNRAMP4 is required for optimal photosynthesis and growth under manganese deficiency. *Plant Physiol.* 152, 1986–1999
58. Burkhead, J.L. *et al.* (2009) Copper homeostasis. *New Phytol.* 182, 799–816
59. Tapken, W. *et al.* (2015) The Clp protease system is required for copper ion-dependent turnover of the PAA2/HMA8 copper transporter in chloroplasts. *New Phytol.* 205, 511–517
60. Shahbaz, M. and Pilon, M. (2019) Conserved Cu-microRNAs in *Arabidopsis thaliana* function in copper economy under deficiency. *Plants (Basel)* 8, E141
61. Tapken, W. *et al.* (2015) Regulation of Cu delivery to chloroplast proteins. *Plant Signal. Behav.* 10, e1046666
62. Solti, A. *et al.* (2012) Uptake and incorporation of iron in sugar beet chloroplasts. *Plant Physiol. Biochem.* 52, 91–97
63. Müller, B. *et al.* (2019) Chloroplasts preferentially take up ferric citrate over iron–nicotianamine complexes in *Brassica napus*. *Planta* 249, 751–763
64. Lopez-Millan, A.F. *et al.* (2016) Chloroplast iron transport proteins – function and impact on plant physiology. *Front. Plant Sci.* 7, 178
65. Shingles, R. *et al.* (2002) Ferrous ion transport across chloroplast inner envelope membranes. *Plant Physiol.* 128, 1022–1030
66. Jeong, J. *et al.* (2008) Chloroplast Fe(III) chelate reductase activity is essential for seedling viability under iron limiting conditions. *Proc. Natl. Acad. Sci. U. S. A.* 105, 10619–10624
67. Solti, A. *et al.* (2014) Functional characterization of the chloroplast ferric chelate oxidoreductase enzyme. *New Phytol.* 202, 920–928
68. Vigan, G. *et al.* (2019) Essential and detrimental – an update on intracellular iron trafficking and homeostasis. *Plant Cell Physiol.* 60, 1420–1439
69. Duy, D. *et al.* (2011) The chloroplast permease PIC1 regulates plant growth and development by directing homeostasis and transport of iron. *Plant Physiol.* 155, 1709–1722
70. Tarantino, D. *et al.* (2011) Identification of an *Arabidopsis* mitoferrinlike carrier protein involved in Fe metabolism. *Plant Physiol. Biochem.* 49, 520–529
71. Conte, S. *et al.* (2009) Multiple antibiotic resistance in *Arabidopsis* is conferred by mutations in a chloroplast-localized transport protein. *Plant Physiol.* 151, 559–573

72. Voith von Voithenberg, L. *et al.* (2019) A novel prokaryote-type ECF/ABC transporter module in chloroplast metal homeostasis. *Front. Plant Sci.* 10, 1264
73. Shimoni-Shor, E. *et al.* (2010) Disruption of Nap14, a plastid-localized non-intrinsic ABC protein in *Arabidopsis thaliana* results in the over-accumulation of transition metals and in aberrant chloroplast structures. *Plant Cell Environ.* 33, 1029–1038
74. Divol, F. *et al.* (2013) The *Arabidopsis* YELLOW STRIPE LIKE4 and 6 transporters control iron release from the chloroplast. *Plant Cell* 25, 1040–1055
75. Zhang, X.Y. *et al.* (2017) *Zea mays* Fe deficiency-related 4 (ZmFDR4) functions as an iron transporter in the plastids of monocots. *Plant J.* 90, 147–163
76. Wang, C. *et al.* (2016) A putative chloroplast-localized $\text{Ca}^{2+}/\text{H}^{+}$ antiporter CCHA1 is involved in calcium and pH homeostasis and required for PSII function in *Arabidopsis*. *Mol. Plant* 9, 1183–1196
77. Krieger-Liszka, A. and Thomine, S. (2018) Importing manganese into the chloroplast: many membranes to cross. *Mol. Plant* 11, 1109–1111
78. Hoecker, N. *et al.* (2017) Plants contain small families of UPF0016 proteins including the PHOTOSYNTHESIS AFFECTED MUTANT71 transporter. *Plant Signal. Behav.* 12, e1278101
79. Frank, J. *et al.* (2019) Chloroplast-localized BICAT proteins shape stromal calcium signals and are required for efficient photosynthesis. *New Phytol.* 221, 866–880
80. Brandenburg, F. *et al.* (2017) The *Synechocystis* manganese exporter Mnx is essential for manganese homeostasis in cyanobacteria. *Plant Physiol.* 173, 1798–1810
81. Gandini, C. *et al.* (2017) The transporter SynPAM71 is located in the plasma membrane and mediates manganese tolerance in *Synechocystis* PCC6803. *New Phytol.* 215, 256–268
82. Abdel-Ghany, S.E. *et al.* (2005) Two P-type ATPases are required for copper delivery in *Arabidopsis thaliana* chloroplasts. *Plant Cell* 17, 1233–1251
83. Catty, P. *et al.* (2011) Biochemical characterization of AtHMA6/PAA1, a chloroplast envelope Cu(I) -ATPase. *J. Biol. Chem.* 286, 36188–36197
84. Sautron, E. *et al.* (2015) HMA6 and HMA8 are two chloroplast Cu^{+} -ATPases with different enzymatic properties. *Biosci. Rep.* 35, e00201
85. Seigneurin-Berny, D. *et al.* (2006) HMA1, a new Cu -ATPase of the chloroplast envelope, is essential for growth under adverse light conditions. *J. Biol. Chem.* 281, 2882–2892
86. Boutigny, S. *et al.* (2014) HMA1 and PAA1, two chloroplast-envelope PIB-ATPases, play distinct roles in chloroplast copper homeostasis. *J. Exp. Bot.* 65, 1529–1540
87. Mikkelsen, M.D. *et al.* (2012) Barley HvHMA1 is a heavy metal pump involved in mobilizing organellar Zn and Cu and plays a role in metal loading into grains. *PLoS One* 7, e49027
88. Kim, Y.Y. *et al.* (2009) AtHMA1 contributes to the detoxification of excess Zn(II) in *Arabidopsis*. *Plant J.* 58, 737–753
89. Palmer, T. and Berks, B.C. (2012) The twin-arginine translocation (Tat) protein export pathway. *Nat. Rev. Microbiol.* 10, 483–496
90. Hamsanathan, S. and Musser, S.M. (2018) The Tat protein transport system: intriguing questions and conundrums. *FEMS Microbiol. Lett.* 365, fny123
91. Ravet, K. *et al.* (2009) Ferritins control interaction between iron homeostasis and oxidative stress in *Arabidopsis*. *Plant J.* 57, 400–412
92. Turowski, V.R. *et al.* (2015) Frataxin is localized to both the chloroplast and mitochondrion and is involved in chloroplast Fe-S protein function in *Arabidopsis*. *PLoS One* 10, e0141443
93. Armas, A.M. *et al.* (2019) Ferredoxin activity of plant frataxin. *Biochimie* 156, 118–122
94. Hu, X. *et al.* (2017) The SUFBC₂D complex is required for the biogenesis of all major classes of plastid Fe-S proteins. *Plant J.* 90, 235–248
95. Nath, K. *et al.* (2016) A nitrogen-fixing subunit essential for accumulating 4Fe-4S-containing photosystem I core proteins. *Plant Physiol.* 172, 2459–2470
96. Molik, S. *et al.* (2001) The Rieske Fe/S protein of the cytochrome *b₆/f* complex in chloroplasts – missing link in the evolution of protein transport pathways in chloroplasts? *J. Biol. Chem.* 276, 42761–42766
97. Roose, J.L. *et al.* (2016) The extrinsic proteins of photosystem II: update. *Planta* 243, 889–908
98. Cao, P. *et al.* (2015) Crystal structure analysis of extrinsic PsbP protein of photosystem II reveals a manganese-induced conformational change. *Mol. Plant* 8, 664–666
99. Bondarava, N. *et al.* (2005) Function of the 23 kDa extrinsic protein of photosystem II as a manganese binding protein and its role in photoactivation. *Biochim. Biophys. Acta* 1708, 63–70
100. Bondarava, N. *et al.* (2007) Manganese binding to the 23 kDa extrinsic protein of photosystem II. *Biochim. Biophys. Acta* 1767, 583–588
101. Blaby-Haas, C.E. *et al.* (2014) Evolution of a plant-specific copper chaperone family for chloroplast copper homeostasis. *Proc. Natl. Acad. Sci. U. S. A.* 111, E5480–E5487
102. Cohn, C.M. *et al.* (2009) Copper delivery by the copper chaperone for chloroplast and cytosolic copper/zinc-superoxide dismutases: regulation and unexpected phenotypes in an *Arabidopsis* mutant. *Mol. Plant* 2, 1336–1350
103. Wei, X. *et al.* (2016) Structure of spinach photosystem II-LHCII supercomplex at 3.2 Å resolution. *Nature* 534, 69–74
104. Su, X. *et al.* (2017) Structure and assembly mechanism of plant C₂S₂M₂-type PSII-LHCII supercomplex. *Science* 357, 815–820
105. van Bezouwen, L.S. *et al.* (2017) Subunit and chlorophyll organization of the plant photosystem II supercomplex. *Nat. Plants* 3, 17080
106. Stroebel, D. *et al.* (2003) An atypical haem in the cytochrome *b₆/f* complex. *Nature* 426, 413–418
107. Hasan, S.S. *et al.* (2013) Quinone-dependent proton transfer pathways in the photosynthetic cytochrome *b₆/f* complex. *Proc. Natl. Acad. Sci. U. S. A.* 110, 4297–4302
108. Pan, X. *et al.* (2018) Structure of the maize photosystem I supercomplex with light-harvesting complexes I and II. *Science* 360, 1109–1113
109. Amunts, A. *et al.* (2010) Structure determination and improved model of plant photosystem I. *J. Biol. Chem.* 285, 3478–3486
110. Mazar, Y. *et al.* (2017) Structure of the plant photosystem I supercomplex at 2.6 Å resolution. *Nat. Plants* 3, 17014
111. Im, S.C. *et al.* (1998) The solution structure of parsley [2Fe-2S] ferredoxin. *Eur. J. Biochem.* 258, 465–477
112. Kurisu, G. *et al.* (2001) Structure of the electron transfer complex between ferredoxin and ferredoxin-NADP⁺ reductase. *Nat. Struct. Biol.* 8, 117–121
113. Moore, J.M. *et al.* (1991) High-resolution solution structure of reduced French bean plastocyanin and comparison with the crystal structure of poplar plastocyanin. *J. Mol. Biol.* 221, 533–555
114. Dietz, K.-J. *et al.* (2016) Redox- and reactive oxygen species-dependent signaling into and out of the photosynthesizing chloroplast. *Plant Physiol.* 171, 1541–1550
115. Leister, D. (2019) Piecing the puzzle together: the central role of reactive oxygen species and redox hubs in chloroplast retrograde signaling. *Antioxid. Redox Signal.* 30, 1206–1219
116. Vigan, G. *et al.* (2013) Signals from chloroplasts and mitochondria for iron homeostasis regulation. *Trends Plant Sci.* 18, 305–311
117. Grillet, L. *et al.* (2018) IRON MAN is a ubiquitous family of peptides that control iron transport in plants. *Nat. Plants* 4, 953–963